

# Simpson-like and Hermite-Hadamard-like Type Inequalities for Harmonically Quasi-convex Functions

Jaekeun Park

Department of Mathematics  
Hanseu University  
Seosan, Chungnam, 356-706, Korea

Copyright © 2014 Jaekeun Park. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

In this paper, by using a generalized integral identity for differentiable functions, the author obtain some new upper bounds of Hermite-Hadamard type inequalities and new Simpson-like type inequalities, for differentiable harmonically quasi-convex functions.

**Mathematics Subject Classification:** 26A51, 26D15

**Keywords:** Hermite-Hadamard type inequality, Simpson type inequality, Hölder's inequality, Harmonically convexity

## 1 Introduction

Many inequalities have been established for convex functions but the most famous is the Hermite-Hadamard's inequality, due to its rich geometrical significance and applications, which is stated as follows: Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a convex function and  $a, b \in I$  with  $a < b$ . Then following double inequalities hold:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}.$$

Hermite-Hadamard's inequalities for convex functions have received renewed attention in recent years and a remarkable variety of refinements and generalizations have been found in [1, 2, 3, 7, 8, 17] and references therein.

Let us recall some definitions of several kinds of convex functions:

**Definition 1.** Let  $I$  be an interval in  $R$ . Then  $f : I \rightarrow R$  is said to be convex on  $I$  if the inequality

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y) \quad (1)$$

holds, for all  $x, y \in I$  and  $t \in [0, 1]$ .

**Definition 2.** Let  $I$  be an interval in  $R_+ = (0, \infty)$ . A function  $f : I \rightarrow R$  is said to be harmonically convex on  $I$  if the inequality

$$f\left(\frac{xy}{tx + (1-t)y}\right) \leq tf(y) + (1-t)f(x) \quad (2)$$

holds, for all  $x, y \in I$  and  $t \in [0, 1]$ . If the inequality in (2) is reversed, then  $f$  is said to be harmonically concave.

In [4], Imdat İşcan established the following result of the Hermite-Hadamard type for harmonically convex functions:

**Theorem 1.1.** Let  $f : I \subseteq R_+ = (0, \infty) \rightarrow R$  be a harmonically convex function on an interval  $I$  and  $f \in L[a, b]$ , where  $a, b \in I$  with  $a < b$ .

$$f\left(\frac{2ab}{a+b}\right) \leq \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \leq \frac{f(a) + f(b)}{2}. \quad (3)$$

Also, in [4, 5], Imdat İşcan established some new Hermite-Hadamard type inequalities, which estimate the difference between the middle and the rightmost terms in (3), for harmonically convex functions:

**Theorem 1.2.** Let  $f : I \subseteq R_+ = (0, \infty) \rightarrow R$  be a differentiable function on the interior  $I^0$  of an interval  $I$  in  $R_+ = (0, \infty)$  and  $f' \in L[a, b]$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|^q$  is harmonically convex function on  $[a, b]$  for  $q \geq 1$ , then the following inequality holds:

$$\begin{aligned} & \left| \frac{f(a) + f(b)}{2} - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\ & \leq \frac{ab(b-a)}{2} \lambda_1^{1-\frac{1}{q}} \left[ \lambda_2 |f'(a)|^q + \lambda_3 |f'(b)|^q \right]^{\frac{1}{q}}, \end{aligned}$$

where

$$\begin{aligned}\lambda_1 &= \frac{1}{ab} - \frac{2}{(b-a)^2} \ln \left( \frac{(a+b)^2}{4ab} \right), \\ \lambda_2 &= -\frac{1}{b(b-a)} + \frac{3a+b}{(b-a)^3} \ln \left( \frac{(a+b)^2}{4ab} \right), \\ \lambda_3 &= \frac{1}{a(b-a)} - \frac{3b+a}{(b-a)^3} \ln \left( \frac{(a+b)^2}{4ab} \right) \\ &= \lambda_1 - \lambda_2.\end{aligned}$$

In this article we consider the following special functions:

**Definition 3.** The hypergeometric function  ${}_2F_1[a, b, c, x]$  is defined for  $|x| < 1$  by the power series

$${}_2F_1[a, b, c, x] = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{x^n}{n!}. \quad (4)$$

Here  $(q)_n$  is the Pochhammer symbol, which is defined by

$$(q)_n = \begin{cases} 1, & n = 0 \\ q(q+1) \cdots (q+n-1), & n > 0. \end{cases}$$

**Definition 4.** The beta function, also called the Euler integral of the first kind, is a special function defined by

$$\beta(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt. \quad (5)$$

In this paper, we give some generalized inequalities connected with the left and right parts of (3), as a result of this, we obtain some generalized Simpson-like and Hermite-Hadamard-like type inequalities for differentiable harmonically quasi-convex functions by using an integral identity for differentiable functions.

## 2 Main results

In order to find some new inequalities of Hermite-Hadamard-like and Simpson-like type inequalities connected with the left and right parts of (3) for functions whose derivatives are harmonically quasi-convex, we need the following lemma [15]:

**Lemma 1.** Let  $f : I \subseteq R_+ = (0, \infty) \rightarrow R$  be a differentiable function on the interior  $I^0$  of an interval  $I$  such that  $f' \in L([a, b])$ , where  $a, b \in I$  with  $a < b$ . Then, for  $h \in (0, 1)$  with  $\frac{1}{n} \leq h \leq \frac{n-1}{n}$  for any  $n \geq 2$  the following identity

$$\begin{aligned} S_a^b(f)(h, n) &= \frac{1}{n} \left[ f(a) + (n-2)f\left(\frac{ab}{A_h(a, b)}\right) + f(b) \right] - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \\ &= ab(a-b) \int_0^1 p(t, h) \frac{1}{A_t^2(a, b)} f'\left(\frac{ab}{A_t(a, b)}\right) dt \end{aligned} \quad (6)$$

holds, for  $t \in [0, 1]$ , where  $A_t(a, b) = (1-t)a + tb$  and

$$p(t, h) = \begin{cases} t - \frac{1}{n} & t \in [0, h] \\ t - \frac{n-1}{n} & t \in (h, 1]. \end{cases}$$

*Proof* By the simple calculation, this is proved.

Now we turn our attention to establish inequalities of Hermit-Hadamard-like and Simpson-like type for differentiable harmonically convex functions.

**Theorem 2.1.** Let  $f : I \subseteq R_+ = (0, \infty) \rightarrow R$  be a differentiable function on  $I^0$ , the interior of an interval  $I$ , such that  $f' \in L([a, b])$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|$  is harmonically quasi-convex on  $[a, b]$ , then, for  $h \in (0, 1)$  with  $\frac{1}{n} \leq h \leq \frac{n-1}{n}$  for any  $n \geq 2$  the following inequality

$$\begin{aligned} &|S_a^b(f)(h, n)| \\ &\leq ab(b-a) \{ \mu_{11}(h, n) + \mu_{12}(h, n) \} \sup \{ |f'(a)|, |f'(b)| \} \end{aligned} \quad (7)$$

holds, where

$$\begin{aligned} \mu_{11}(h, n) &= \frac{(2-h-nh)a + hb}{na(b-a)A_h(a, b)} + \frac{1}{(b-a)^2} \ln \left[ \frac{aA_h(a, b)}{A_{\frac{1}{n}}^2(a, b)} \right], \\ \mu_{12}(h, n) &= -\frac{(1-h)a + (1+h-n+nh)b}{nb(b-a)A_h(a, b)} \\ &\quad + \frac{1}{(b-a)^2} \ln \left[ \frac{bA_h(a, b)}{A_{\frac{1}{n}}^2(a, b)} \right]. \end{aligned}$$

*Proof* From Lemma 1, since  $|f'|$  is harmonically convex on  $[a, b]$ , we have

$$\begin{aligned} &|S_a^b(f)(h, n)| \\ &\leq ab(b-a) \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} \left| f'\left(\frac{ab}{A_t(a, b)}\right) \right| dt \\ &\quad + \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} \left| f'\left(\frac{ab}{A_t(a, b)}\right) \right| dt \\ &\leq \left\{ \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} dt + \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} dt \right\} \sup \{ |f'(a)|, |f'(b)| \}. \end{aligned} \quad (8)$$

Note that

$$\begin{aligned} (a) \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} dt \\ = \frac{(2 - h - nh)a + hb}{na(b - a)A_h(a, b)} + \frac{1}{(b - a)^2} \ln \left[ \frac{aA_h(a, b)}{A_{\frac{1}{n}}^2(a, b)} \right], \end{aligned} \quad (9)$$

$$\begin{aligned} (b) \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} dt \\ = -\frac{(1 - h)a + (1 + h - n + nh)b}{nb(b - a)A_h(a, b)} \\ + \frac{1}{(b - a)^2} \ln \left[ \frac{bA_h(a, b)}{A_{\frac{1}{n}}^2(a, b)} \right]. \end{aligned} \quad (10)$$

By substituting (9) and (10) in (8), we get the desired result.

**Corollary 2.1.** *In Theorem 2.1, one has:*

$$\begin{aligned} |S_a^b(f)(\tfrac{1}{2}, 6)| \\ \leq ab(b - a) \{ \mu_{11}(\tfrac{1}{2}, 6) + \mu_{12}(\tfrac{1}{2}, 6) \} \sup \{ |f'(a)|, |f'(b)| \}. \end{aligned}$$

**Theorem 2.2.** *Let  $f : I \subseteq R_+ = (0, \infty) \rightarrow R$  be a differentiable function on the interior  $I^0$  of an interval  $I$  and  $f' \in L([a, b])$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|^q$  is harmonically quasi-convex on  $[a, b]$  for  $q > 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$ , then, for  $h \in (0, 1)$  with  $\frac{1}{n} \leq h \leq \frac{n-1}{n}$  for any  $n \geq 2$  the following inequality*

$$\begin{aligned} |S_a^b(f)(h, n)| \\ \leq ab(b - a) \left\{ (\mu_{21}(h, n) + \mu_{22}(h, n))^{\frac{1}{p}} h^{\frac{1}{q}} \right. \\ \left. + (\mu_{23}(h, n) + \mu_{24}(h, n))^{\frac{1}{p}} (1 - h)^{\frac{1}{q}} \right\} \\ \times \left( \sup \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}}. \end{aligned} \quad (11)$$

holds, where

$$\begin{aligned}
 \mu_{21}(h, n) &= \frac{1}{(1+p)n^{1+p}a^{2p}} {}_2F_1\left[1, 2p, 2+p, -\frac{b-a}{na}\right], \\
 \mu_{22}(h, n) &= A_{\frac{1}{n}}^{1-2p}(a, b) \frac{n^p}{(a-b)^{1+p}} \beta(1-2p, 1+p) \\
 &\quad + \frac{A_h^{1-2p}(a, b) A_{\frac{1}{n}}^p(a, b)}{(2p-1)(a-b)^{1+p}} {}_2F_1\left[1-2p, -p, 2-2p, \frac{A_h(a, b)}{A_{\frac{1}{n}}(a, b)}\right], \\
 \mu_{23}(h, n) &= A_{\frac{1}{n}}^{1-p}(b, a) \frac{\beta(1-2p, 1+p)}{(b-a)^{1+p}} \\
 &\quad + \frac{A_h^{1-2p}(a, b) A_{\frac{1}{n}}^p(b, a)}{2p-1} \frac{A_{\frac{1}{n}}^p(b, a)}{(b-a)^{1+p}} {}_2F_1\left[1-2p, -p, 2-2p, \frac{A_h(a, b)}{A_{\frac{1}{n}}(b, a)}\right], \\
 \mu_{24}(h, n) &= A_{\frac{1}{n}}^{1-p}(b, a) \frac{\beta(1-2p, 1+p)}{(a-b)^{1+p}} \\
 &\quad + \frac{b^{1-2p} A_{\frac{1}{n}}^p(b, a)}{(2p-1)(a-b)^{1+p}} {}_2F_1\left[1-2p, -p, 2-2p, \frac{b}{A_{\frac{1}{n}}(b, a)}\right].
 \end{aligned}$$

*Proof* From Lemma 1 and by the Hölder integral inequality, we have

$$\begin{aligned}
 &|S_a^b(f)(h, n)| \\
 &\leq ab(b-a) \left[ \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right| dt \right. \\
 &\quad \left. + \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right| dt \right] \\
 &\leq ab(b-a) \left[ \left\{ \int_0^h \frac{|t - \frac{1}{n}|^p}{A_t^{2p}(a, b)} dt \right\}^{\frac{1}{p}} \left\{ \int_0^h \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \right\}^{\frac{1}{q}} \right. \\
 &\quad \left. + \left\{ \int_h^1 \frac{|t - \frac{n-1}{n}|^p}{A_t^{2p}(a, b)} dt \right\}^{\frac{1}{p}} \left\{ \int_h^1 \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \right\}^{\frac{1}{q}} \right]. \tag{12}
 \end{aligned}$$

Note that

$$\begin{aligned}
 (a) \quad &\int_0^h \frac{|t - \frac{1}{n}|^p}{A_t^{2p}(a, b)} dt \\
 &= \int_0^{\frac{1}{n}} \frac{(\frac{1}{n} - t)^p}{A_t^{2p}(a, b)} dt + \int_{\frac{1}{n}}^h \frac{(t - \frac{1}{n})^p}{A_t^{2p}(a, b)} dt \\
 &= \mu_{21}(h, n) + \mu_{22}(h, n), \tag{13}
 \end{aligned}$$

$$\begin{aligned}
(b) \int_h^1 \frac{\left|t - \frac{n-1}{n}\right|^p}{A_t^{2p}(a, b)} dt \\
= \int_h^{\frac{n-1}{n}} \frac{\left(\frac{n-1}{n} - t\right)^p}{A_t^{2p}(a, b)} dt + \int_{\frac{n-1}{n}}^1 \frac{\left(t - \frac{n-1}{n}\right)^p}{A_t^{2p}(a, b)} dt \\
= \mu_{23}(h, n) + \mu_{24}(h, n).
\end{aligned} \tag{14}$$

Since  $|f'|^q$  is harmonically quasi-convex on  $[a, b]$  for  $q > 1$ , we have

$$\begin{aligned}
(i) \int_0^h \left|f'\left(\frac{ab}{A_t(a, b)}\right)\right|^q dt \\
\leq \int_0^h \sup \{|f'(a)|^q, |f'(b)|^q\} dt \\
= h \sup \{|f'(a)|^q, |f'(b)|^q\},
\end{aligned} \tag{15}$$

$$\begin{aligned}
(ii) \int_h^1 \left|f'\left(\frac{ab}{A_t(a, b)}\right)\right|^q dt \\
\leq \int_h^1 \sup \{|f'(a)|^q, |f'(b)|^q\} dt \\
= (1 - h) \sup \{|f'(a)|^q, |f'(b)|^q\}.
\end{aligned} \tag{16}$$

By substituting (13)-(16) in (12), we get the desired result (11).

**Theorem 2.3.** Let  $f : I \subseteq R_+ = (0, \infty) \rightarrow R$  be a differentiable function on the interior  $I^0$  of an interval  $I$  and  $f' \in L([a, b])$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|^q$  is harmonically quasi-convex on  $[a, b]$  for  $q > 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$ , then, for  $h \in (0, 1)$  with  $\frac{1}{n} \leq h \leq \frac{n-1}{n}$  for any  $n \geq 2$  the following inequality

$$\begin{aligned}
& \left|S_a^b(f)(h, n)\right| \\
& \leq ab(b-a) \left\{ \left(\frac{1+(nh-1)^{1+p}}{(1+p)n^{1+p}}\right)^{\frac{1}{p}} \mu_{31}^{\frac{1}{q}}(h) \right. \\
& \quad \left. + \left(\frac{1+(n-nh-1)^{1+p}}{(1+p)n^{1+p}}\right)^{\frac{1}{p}} \mu_{32}^{\frac{1}{q}}(h) \right\} \\
& \quad \times \left\{ \sup \{|f'(a)|^q, |f'(b)|^q\} \right\}^{\frac{1}{q}}
\end{aligned} \tag{17}$$

holds, where

$$\begin{aligned}
\mu_{31}(h) &= \frac{A_h^{1-2q}(a, b) - a^{1-2q}}{(1-2q)(b-a)}, \\
\mu_{32}(h) &= \frac{b^{1-2q} - A_h^{1-2q}(a, b)}{(1-2q)(b-a)}.
\end{aligned}$$

*Proof* From Lemma 1 and by the Hölder integral inequality, we have

$$\begin{aligned}
 & |S_a^b(f)(h, n)| \\
 & \leq ab(b-a) \left[ \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right| dt \right. \\
 & \quad \left. + \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right| dt \right] \\
 & \leq ab(b-a) \left[ \left\{ \int_0^h \left| t - \frac{1}{n} \right|^p dt \right\}^{\frac{1}{p}} \right. \\
 & \quad \times \left\{ \int_0^h \frac{1}{A_t^{2q}(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \right\}^{\frac{1}{q}} \\
 & \quad + \left\{ \int_h^1 \left| t - \frac{n-1}{n} \right|^p dt \right\}^{\frac{1}{p}} \\
 & \quad \times \left\{ \int_h^1 \frac{1}{A_t^{2q}(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \right\}^{\frac{1}{q}} \Big]. \tag{18}
 \end{aligned}$$

Note that

$$\int_0^h \left| t - \frac{1}{n} \right|^p dt = \frac{1 + (nh - 1)^{1+p}}{(1+p)n^{1+p}}, \tag{19}$$

$$\int_h^1 \left| t - \frac{n-1}{n} \right|^p dt = \frac{1 + (n - nh - 1)^{1+p}}{(1+p)n^{1+p}}. \tag{20}$$

Since  $|f'|^q$  is harmonically quasi-convex on  $[a, b]$  for  $q > 1$ , we have

$$\begin{aligned}
 (i) \quad & \int_0^h \frac{1}{A_t^{2q}(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \\
 & \leq \left( \int_0^h \frac{1}{A_t^{2q}(a, b)} dt \right) \sup \{ |f'(a)|^q, |f'(b)|^q \} \\
 & = \mu_{31}(h) \sup \{ |f'(a)|^q, |f'(b)|^q \}, \tag{21}
 \end{aligned}$$

$$\begin{aligned}
 (ii) \quad & \int_h^1 \frac{1}{A_t^{2q}(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \\
 & \leq \left( \int_h^1 \frac{1}{A_t^{2q}(a, b)} dt \right) \sup \{ |f'(a)|^q, |f'(b)|^q \} \\
 & = \mu_{32}(h) \sup \{ |f'(a)|^q, |f'(b)|^q \}. \tag{22}
 \end{aligned}$$

By substituting (19)-(22) in (18), we get the desired result (17).



**Corollary 2.2.** *In Theorem 2.3, one has:*

$$\begin{aligned} |S_a^b(f)(\tfrac{1}{2}, 6)| &\leq ab(b-a) \left\{ \frac{1+2^{1+p}}{(1+p)6^{1+p}} \right\}^{\frac{1}{p}} \left\{ \mu_{31}^{\frac{1}{q}}(\tfrac{1}{2}) + \mu_{32}^{\frac{1}{q}}(\tfrac{1}{2}) \right\} \\ &\quad \times \left\{ \sup \{ |f'(a)|^q, |f'(b)|^q \} \right\}^{\frac{1}{q}}. \end{aligned} \quad (23)$$

**Theorem 2.4.** *Let  $f : I \subseteq R_+ = (0, \infty) \rightarrow R$  be a differentiable function on the interior  $I^0$  of an interval  $I$  and  $f' \in L([a, b])$ , where  $a, b \in I$  with  $a < b$ . If  $|f'|^q$  is harmonically quasi-convex on  $[a, b]$  for  $q \geq 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$ , then, for  $h \in (0, 1)$  with  $\frac{1}{n} \leq h \leq \frac{n-1}{n}$  for any  $n \geq 2$  the following inequality*

$$\begin{aligned} |S_a^b(f)(h, n)| &\leq ab(b-a) \left( \mu_{11}(h, n) + \mu_{12}(h, n) \right) \\ &\quad \times \left( \sup \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}}. \end{aligned} \quad (24)$$

holds, where  $\mu_{1i}(\cdot, \cdot)$  ( $i = 1, 2$ ) are defined in Theorem 2.1.

*Proof* From Lemma 1 and by the power mean integral inequality, we have

$$\begin{aligned} |S_a^b(f)(h, n)| &\leq ab(a-b) \left[ \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right| dt \right. \\ &\quad \left. + \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right| dt \right] \\ &\leq ab(a-b) \left[ \left\{ \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} dt \right\}^{\frac{1}{p}} \right. \\ &\quad \times \left\{ \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \right\}^{\frac{1}{q}} \\ &\quad + \left\{ \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} dt \right\}^{\frac{1}{p}} \\ &\quad \times \left. \left\{ \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \right\}^{\frac{1}{q}} \right]. \end{aligned} \quad (25)$$

Since  $|f'|^q$  is harmonically quasi-convex on  $[a, b]$  for  $q \geq 1$ , we have

$$\begin{aligned} (i) \quad & \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \\ & \leq \left( \int_0^h \frac{|t - \frac{1}{n}|}{A_t^2(a, b)} dt \right) \sup \{ |f'(a)|^q, |f'(b)|^q \} \\ & = \mu_{11}(h, n) \sup \{ |f'(a)|^q, |f'(b)|^q \}, \end{aligned} \quad (26)$$

$$\begin{aligned} (ii) \quad & \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} \left| f' \left( \frac{ab}{A_t(a, b)} \right) \right|^q dt \\ & \leq \left( \int_h^1 \frac{|t - \frac{n-1}{n}|}{A_t^2(a, b)} dt \right) \sup \{ |f'(a)|^q, |f'(b)|^q \} \\ & = \mu_{12}(h, n) \sup \{ |f'(a)|^q, |f'(b)|^q \}. \end{aligned} \quad (27)$$

By substituting (9)-(10) and (26)-(27) in (25), we get the desired result (24).

**Corollary 2.3.** *In Theorem 2.3, one has:*

$$\begin{aligned} & |S_a^b(f)(\tfrac{1}{2}, 6)| \\ & \leq ab(b-a) \left( \mu_{11}(\tfrac{1}{2}, 6) + \mu_{12}(\tfrac{1}{2}, 6) \right) \\ & \quad \times \left( \sup \{ |f'(a)|^q, |f'(b)|^q \} \right)^{\frac{1}{q}}. \end{aligned} \quad (28)$$

## References

- [1] S. S. Dragomir, Hermite-Hadamard's type inequalities for convex functions of selfadjoint operators in Hilbert spaces, *Linear Algebra Appl.* **436** (5) (2012), 1503–1515. <http://dx.doi.org/10.1016/j.laa.2011.08.050>
- [2] S. S. Dragomir, Superadditivity and monotonicity of some functionals associated with the Hermite-Hadamard inequality for convex functions in linear spaces, *Rocky Mountain J. Math.* **42**(5) (2012), 1447–1459. <http://dx.doi.org/10.1216/RMJ-2012-42-5-1447>
- [3] V. N. Huy, N. T. Chung, Some generalizations of the Fejér and Hermite-Hadamard inequalities in Hölder spaces, *J. Appl. Math. Inform.* **29**(3-4) (2011), 859–868.
- [4] İmdat İşcan, Hermite-Hadamard type inequalities for harmonically convex functions, *arXiv:1303.6089v1[math.CA]25 Mar 2013* (2013).

- [5] İmdat İşcan, Hermite-Hadamard and Simpson-like type inequalities for differentiable harmonically convex functions, presented.
- [6] A. P. Ji, T. Y. Zhang, F. Qi, Integral Inequalities of Hermite-Hadamard Type for  $(\alpha, m)$ -GA-Convex Functions, *Journal of Function Spaces and Applications*, **2013** (2013), Article ID 823856, 8 pages. <http://dx.doi.org/10.1155/2013/823856>
- [7] M. Klaričić Bakula, J. Pečarić, Note on some Hadamard-type inequalities, *JIPAM. J. Inequal. Pure Appl. Math.* **5(3)** (2004), Article 74, 9 pp.
- [8] M. Klaričić Bakula, M. E. Özdemir, J. Pečarić, Hadamard type inequalities for  $m$ -convex and  $(\alpha, m)$ -convex functions, *JIPAM. J. Inequal. Pure Appl. Math.* **9(4)** (2008), Article 96, 12 pp.
- [9] Meihui Qu, Wenjun Liu, J. Park, Some new Hermite-Hadamard-type inequalities for geometric-arithmetically  $s$ -convex functions, Presented
- [10] V. G. Miheşan, "A generalization of the convexity" in Seminar on Functional Equations, *Approx. and Convex.*, Cluj-Napoca, Romania, 1993.
- [11] C. P. Niculescu, Convexity according to the geometric mean, *Math. Inequal. Appl.*, **3(2)** (2000), 155–167. <http://dx.doi.org/10.7153/mia-03-19>
- [12] C. P. Niculescu, Convexity according to means, *Math. Inequal. Appl.*, **6(4)** (2003), 571–579. <http://dx.doi.org/10.7153/mia-06-53>
- [13] M. E. Özdemir, A. Ekinici, A. O. Akdemir, Generalizations of integral inequalities for functions whose second derivatives are convex and  $m$ -convex, *Miskolc Mathematical Notes*, **13(2)** (2012), 441–457.
- [14] J. Park, Some new Hermite-Hadamard-type inequalities for twice differentiable geometric-arithmetically  $s$ -convex functions, *Appl. Math. Sci.*, **7(87)** (2013), 4309–4325. <http://dx.doi.org/10.12988/ams.2013.36303>
- [15] J. Park, Generalization of some Simpson-type inequalities via differentiable  $s$ -convex mappings in the second sense, *Inter. Journ. of Math. and Math Sciences*, **2011** (2011), Art ID:493531. <http://dx.doi.org/10.1155/2011/493531>.
- [16] Feng Qi, Bo-Yan Xi, Some integral inequalities of Simpson type for GA- $\varepsilon$ -convex functions, *Georgian Math. J.*, **20(4)** (2013), 775–788. <http://dx.doi.org/10.1515/gmj-2013-0043>
- [17] M. Z. Sarikaya, On new Hermite Hadamard Fejér type integral inequalities, *Stud. Univ. Babeş-Bolyai Math.* **57(3)** (2012), 377–386.

- [18] T. Y. Zhang, A. P. Ji, F. Qi, Some inequalities of Hermite-Hadamard type for GA-Convex functions with applications to means. *Le Matematiche*, **68(1)** (2013), 229–239.

**Received: June 5, 2014**